Author Index

Aguilar, M. I., see Hearn, M. T. W. 33

Airaudo, Ch. B.
——, Gayte-Sorbier, A. and Creusevau, R.
Thin-Layer chromatographic analysis of phenolic age resisters in elastomers in contact with the body, drugs and foods 407

Ali, I., see Bhushan, R. 460

Aubel, M.
—— and Rogers, L. B.
Effects of pretreatment on the enantioselectivity of silica-bound bovine serum albumin used as high-performance liquid chromatographic stationary phases 415

Aue, W. A., see Siu, K. W. M. 143

Baba, S.
——, Suzuki, Y., Sasaki, Y. and Horie, M.
Further study of the synchronized accumulating radioisotope detector for high-performance liquid chromatography 157

Baker, G. B.
——, Wong, J. T. F., Coutts, R. T. and Pasutto, F. M.
Simultaneous extraction and quantitation of several bioactive amines in cheese and chocolate 317

Batley, G. E., see Low, G. K.-C. 199

Bhushan, R.
—— and Ali, I.
Resolution of enantiomeric mixtures of phenylthiohydantoin amino acids on (+)-tartaric acid-impregnated silica gel plates 460

Biggs, W. R., see Jinno, K. 75

Björkholm, E., see Rudling, J. 239

Blank, M. L.
——, Cress, E. A. and Snyder, F.
Separation and quantitation of phospholipid subclasses as their diradylglycerobenzoate derivatives by normal-phase high-performance liquid chromatography 421

Bossi, M. L., see Righetti, P. G. 123

Brieva, A., see Rubio, N. 447

Brockbank, C. I., see Low, G. K.-C. 199

Chapin, F., see Folena-Wasserman, G. 225

Chatkittikunrong, W., see Onodera, S. 295

Chaves Das Neves, H. J.
—— and Vasconcelos, A. M. P.
Capillary gas chromatography of amino acids, including asparagine and glutamine: sensitive gas chromatographic–mass spectrometric and selected ion monitoring gas chromatographic–mass spectrometric detection of the N,O(S)-*tert.*-butyldimethylsilyl derivatives 249

Chiavari, G.
——, Vitali, P. and Galletti, G. C.
Electrochemical detection in the high-performance liquid chromatography of polyphenols (vegetables tannins) 426

Christie, W. W.
—— and Stefanov, K.
Separation of picolinyl ester derivatives of fatty acids by high-performance liquid chromatography for identification by mass spectrometry 259

Cohen, A. I., see Jemal, M. 442

Coutts, R. T., see Baker, G. B. 317

Crescentini, G.
——, Mangani, F., Mastrogiacomo, A. R. and Palma, P.
Discussion of thermodynamic data obtained by adsorption gas chromatography of hydrocarbons on graphitized carbon blacks 83

Cress, E. A., see Blank, M. L. 421

Creusevau, R., see Airaudo, Ch. B. 407

Das Neves, H. J., Chaves, see Chaves Das Neves, H. J. 249

De Vos, F. L.
——, Robertson, D. M. and Hearn, M. T. W.
Effect of mass loadability, protein concentration and *n*-alkyl chain length on the reversed-phase high-performance liquid chromatographic behaviour of bovine serum albumin and bovine follicular fluid inhibin 17

Di Nunzio, C.
——, Parisi, G., Santoro, P. and Ricci, P. A.
Determination of phenols and cresols in aluminium-backed paper by high-performance liquid chromatography 454

Doehl, J.
——, Farbrot, A., Greibrokk, T. and Iversen, B.
Supercritical fluid chromatographic separation of low- to medium-polarity compounds that are difficult to elute from packed columns 175

Donnelly, J. R.
——, Munslow, W. D., Mitchum, R. K. and Sovocool, G. W.
Correlation of structure with retention index for chlorinated dibenzo-*p*-dioxins 51

Farbrot, A., see Doehl, J. 175

Ferenczi-Fodor, K.
——, Kovács, I. and Szepesi, G.
Separation and determination of steroid isomers on amino-bonded silica by conventional and overpressurized thin-layer chromatography 464

Fetzer, J. C., see Jinno, K. 75
Folena-Wasserman, G.
——, Sitrin, R. D., Chapin, F. and Snader, K. M.
Affinity chromatography of glycopeptide antibiotics 225
Frisvad, J. C.
High-performance liquid chromatographic determination of profiles of mycotoxins and other secondary metabolites 333
Gaget, C., see Wagner, J. 211
Galletti, G. C., see Chiavari, G. 426
Gasanova, G. Z., see Zaslavsky, B. Yu. 95
Gayte-Sorbier, A., see Airaudo, Ch. B. 407
Gelfi, C., see Righetti, P. G. 123
Greibrokk, T., see Doehl, J. 175
Haddad, P. R.
—— and Sekulic, S.
Computer routine for selection of the mobile phase search area in optimisation methods 65
Hearn, M. T. W.
—— and Aguilar, M. I.
High-performance liquid chromatography of amino acids, peptides and proteins. LXIX. Evaluation of retention and bandwidth relationships of myosin-related peptides separated by gradient elution reversed-phase high-performance liquid chromatography 33
——, see De Vos, F. L. 17
Heintzelmann, B., see Wagner, J. 211
Horie, M., see Baba, S. 157
Hunchak, K.
—— and Suffet, I. H.
Analysis of acetone–hexane artifacts produced in the Soxhlet extraction of solid environmental samples 185
Hurst, W. J.
——, Snyder, K. P. and Martin, Jr., R. A.
High-performance liquid chromatographic determination of the mycotoxins patulin, penicillic acid, zearalenone and sterigmatocystin in artificially contaminated cocoa beans 389
Hurtubise, R. J., see Ruckmick, S. C. 277
Iversen, B., see Doehl, J. 175
Iwinski, G.
—— and Jenke, D. R.
Determination of alcohols in pharmaceuticals by ion-exclusion chromatography 397
Jacobs, W. A.
o-Phthalaldehyde–sulfite derivatization of primary amines for liquid chromatography–electrochemistry 435
Jemal, M.
—— and Cohen, A. I.
Enantiomeric purity determination of L-proline benzyl ester by chiral column gas chromatography 442
Jenke, D. R., see Iwinski, G. 397
Jinno, K.
——, Nagoshi, T., Tanaka, N., Okamoto, M., Fetzer, J. C. and Biggs, W. R.
Elution behaviour of planar and non-planar polycyclic aromatic hydrocarbons on various chemically bonded stationary phases in liquid chromatography 75
Jones, K.
Optimisation procedure for the silanisation of silicas for reversed-phase high-performance liquid chromatography. I. Elimination of non-significant variables 1
——
Optimisation procedure for the silanisation of silicas for reversed-phase high-performance liquid chromatography. II. Detailed examination of significant variables 11
Kahle, V., see Krejči, M. 133
Kai, M., see Ohno, M. 309
Kiba, N., see Olesik, S. V. 165
Klesper, E., see Leyendecker, D. 101
Kovács, I., see Ferenczi-Fodor, K. 464
Krejči, M.
—— and Kahle, V.
New principle of sample introduction integrated with mobile phase delivery for microcolumn liquid chromatography 133
Leyendecker, D.
——, Leyendecker, D., Schmitz, F. P. and Klesper, E.
Influence of temperature, pressure and eluent composition on the behaviour of binary eluents under sub- and supercritical conditions. Pentane and 1,4-dioxane 101
——, see Leyendecker, D. 101
Low, G. K.-C.
——, Batley, G. E. and Brockbank, C. I.
Solvent-induced photodegradation as a source of error in the analysis of polycyclic aromatic hydrocarbons 199
Mahmudov, A. U., see Zaslavsky, B. Yu. 95
Mangani, F., see Crescentini, G. 83
Martin, Jr., R. A., see Hurst, W. J. 389
Mastrogiacomo, A. R., see Crescentini, G. 83
Miheeva, L. M., see Zaslavsky, B. Yu. 95
Miller, J. H. McB.
——, Pascal, C. and Tissieres, M.
Development of a reversed-phase ion-pairing liquid chromatographic assay of daunorubicin for pharmacopoeial use 361
Mitchum, R. K., see Donnelly, J. R. 51
Munslow, M. D., see Donnelly, J. R. 51
Nagoshi, T., see Jinno, K. 75
Neves, H. J. Chaves Das, see Chaves Das Neves, H. J. 249

Nishikawa, Y.
Liquid chromatographic determination of aliphatic diamines in water via derivatization with acetylacetone 349
Novotný, M. V., see Olesik, S. V. 165
Nunzio, C. Di, see Di Nunzio, C. 454
Ogawa, M., see Onodera, S. 267
Ohkura, Y., see Ohno, M. 309
Ohno, M.
——, Kai, M. and Ohkura, Y.
On-line post-column fluorescence derivatization of arginine-containing peptides in high-performance liquid chromatography 309
Okamoto, M., see Jinno, K. 75
Olesik, S. V.
——, Steger, J. L., Kiba, N., Roth, M. and Novotný, M. V.
High-precision apparatus for physico-chemical measurements by capillary supercritical fluid chromatography 165
Onodera, S.
——, Chatkittikunrong, W., Saito, K., Phongbetchara, R. and Tabucanon, M.
Characterization and determination of lipophilic hydrocarbons in the Chao Phraya, Bang Pakong and Tha-Chin rivers and the Upper Gulf of Thailand 295
——, Ogawa, M. and Suzuki, S.
Chemical changes of organic compounds in chlorinated water. XIII. Gas chromatographic–mass spectrometric studies of the reactions of Irgasan DP 300 [5-chloro-2-(2,4-dichlorophenoxy)-phenol] with chlorine in dilute aqueous solution 267
Palma, P., see Crescentini, G. 83
Parisi, G., see Di Nunzio, C. 454
Pascal, C., see Miller, J. H. McB. 361
Pasutto, F. M., see Baker, G. B. 317
Phongbetchara, R., see Onodera, S. 295
Ricci, P. A., see Di Nunzio, C. 454
Righetti, P. G.
——, Gelfi, C. and Bossi, M. L.
Hydrophobic interaction between alkaline Immobilines and ferritin during isoelectric focusing in immobilized pH gradients 123
Robertson, D. M., see De Vos, F. L. 17
Rogers, L. B., see Aubel, M. 415
Roth, M., see Olesik, S. V. 165
Rubio, M., see Rubio, N. 447
Rubio, N.
——, Brieva, A. and Rubio, M.
Purification of allergens by high-performance liquid chromatography. II. Purification of the major allergen of the mite *Dermatophagoides pteronyssinus* (P_1) 447
Ruckmick, S. C.
——, Hurtubise, R. J. and Silver, H. F.
Separation and characterization of nitrogen heterocycle and hydroxyl aromatic compounds in non-distillable coal-derived liquids 277
Rudling, J.
—— and Björkholm, E.
Irreversibility effects in liquid desorption of organic solvents from activated carbon 239
Rudolph, M.
—— and Steinhart, H.
Determination of Carazolol in tissues of pigs by high-performance liquid chromatography 371
Saito, K., see Onodera, S. 295
Salagoity-Auguste, M.-H.
——, Tricard, C. and Sudraud, P.
Dosage simultané des aldéhydes aromatiques et des coumarines par chromatographie liquide haute performance. Application aux vins et eaux-de-vie vieillis en fût de chêne 379
Santoro, P., see Di Nunzio, C. 454
Sasaki, Y., see Baba, S. 157
Schmitz, F. P., see Leyendecker, D. 101
Sekulic, S., see Haddad, P. R. 65
Silver, H. F., see Ruckmick, S. C. 277
Sitrin, R. D., see Folena-Wasserman, G. 225
Siu, K. W. M.
—— and Aue, W. A.
Steady-state impedance and electron distribution in a ^{63}Ni electron-capture detector 143
Snader, K. M., see Folena-Wasserman, G. 225
Snyder, F., see Blank, M. L. 421
Snyder, K. P., see Hurst, W. J. 389
Sovocool, G. W., see Donnelly, J. R. 51
Stefanov, K., see Christie, W. W. 259
Steger, J. L., see Olesik, S. V. 165
Steinhart, H., see Rudolph, M. 371
Sudraud, P., see Salagoity-Auguste, M.-H. 379
Suffet, I. H., see Hunchak, K. 185
Suzuki, S., see Onodera, S. 267
Suzuki, Y., see Baba, S. 157
Szepesi, G., see Ferenczi-Fodor, K. 464
Tabucanon, M., see Onodera, S. 295
Tanaka, N., see Jinno, K. 75
Tissieres, M., see Miller, J. H. McB. 361
Tricard, C., see Salagoity-Auguste, M.-H. 379
Vasconcelos, A. M. P., see Chaves Das Neves, H. J. 249
Vitali, P., see Chiavari, G. 426
Vos, F. L., de, see De Vos, F. L. 17

Wagner, J.
——, Wolf, E., Heintzelmann, B. and Gaget, C.
Chiral separation of enantiomers of substituted α- and β-alanine and γ-aminobutyric acid analogues by gas chromatography and high-performance liquid chromatography 211
Wolf, E., see Wagner, J. 211
Wong, J. T. F., see Baker, G. B. 317
Zaslavsky, B. Yu.
——, Miheeva, L. M., Gasanova, G. Z. and Mahmudov, A. U.
Influence of inorganic electrolytes on partitioning of non-ionic solutes in an aqueous dextran–poly(ethylene glycol) biphasic system 95